BIO Web of Conferences **149**, 01011 (2024) https://doi.org/10.1051/bioconf/202414901011
Genetic Resources 2024

Comparative analysis of cerebral circulation impairment in humans and horses: implications for veterinary and human medicine

N. Khaidarov[1], *M.* Abdullayeva[1*], and *M.* Tursunova[1]

[1]Tashkent State Dental Institute, Tashkent, Uzbekistan

Abstract: The comparison of cerebral circulation impairment in humans and horses provides important insights into the dynamics of blood flow in the brain across different species, connecting veterinary and human medicine. This analysis brings together findings from various studies to highlight the similarities in anatomy and physiology between humans and horses, which can improve our understanding of cerebrovascular health. Research emphasizes the importance of maintaining stable cerebral blood flow to prevent damage from lack of oxygen, which is essential for the brain's metabolism in both humans and horses. Studies on the factors affecting overall blood flow in the body reveal strong connections between cardiovascular health and blood flow to the brain, showing how baroreflex sensitivity and blood pressure play a role in maintaining brain function. Understanding the reactivity of blood vessels in the brain is associated with neurological disorders, emphasizing the need for non-invasive evaluation techniques that can be used for both humans and horses. Research on the coupling of nerve and blood vessel activity further explains how blood flow is regulated during brain activity, highlighting potential weaknesses under unhealthy conditions. While the role of exercise in promoting cerebrovascular health is recognized, there are still gaps in our knowledge of its effects on the brain, especially in horses. Advanced imaging techniques like MRI are presented as essential tools for assessing the health of small blood vessels and blood flow in the brain, offering potential applications in the early detection and treatment of cerebrovascular diseases. By comparing the dynamics of blood flow in the brain in humans and horses, this review not only increases our understanding but also suggests a more integrated approach to diagnosis and treatment, improving practices in both fields. These insights emphasize the common principles and specific differences in cerebrovascular health in different species, and support the need for collaborative research in veterinary and human medicine.

1 Introduction

The comparative analysis of cerebral circulation impairment in both humans and horses presents a unique intersection of veterinary and human medicine, emphasizing the

[*] Corresponding author: 4research2023@gmail.com

significance of understanding cerebrovascular dynamics across species. The literature provides a foundational framework for this analysis, beginning with [1], who systematically reviewed the effects of anesthetics on cerebral perfusion and oxygenation in pigs. Their findings highlight the anatomical and physiological similarities between the porcine and human brains, particularly regarding cerebral blood flow (CBF) regulation and the intrinsic autoregulatory mechanisms that safeguard optimal cerebral metabolism. This study underscores the necessity of maintaining stable CBF to prevent ischemic damage, a principle that resonates across species, including horses.

In [2], the authors delve into the anatomical aspects of cerebrovascular disease, offering insights into the classical stroke syndromes and the microcirculatory anatomy that underpins cerebrovascular health. Their exploration of the blood-brain barrier (BBB) components and the compensatory capacities of venous structures against thrombosis provides a critical perspective on how anatomical variations may influence cerebrovascular pathology in both humans and horses.

[3] further contributes to this discourse by investigating systemic hemodynamic factors and their impacts on cerebral perfusion. The study establishes a significant relationship between baroreflex sensitivity and regional cerebral perfusion, particularly in the hippocampus, linking systemic cardiovascular health to cerebral function. This connection is crucial for understanding how similar physiological mechanisms may manifest in both humans and horses, especially in the context of exercise and its effects on vascular health.

[4] address cerebrovascular reactivity (CVR) and its implications for cognitive function, revealing that impaired CVR is associated with various neurological disorders. Their systematic review emphasizes the importance of assessing CVR through non-invasive methods, a concept that can be extended to veterinary practices in evaluating equine neurological health. The implications of their findings suggest that both species may share vulnerabilities in cerebrovascular function that warrant further exploration.

[5] provide an integrative overview of cerebrovascular function, discussing neurovascular coupling and the mechanisms that regulate blood flow in response to neuronal activity. Their insights into how these mechanisms adapt under pathological conditions highlight the critical importance of understanding cerebrovascular health in both human and equine populations, particularly in relation to cognitive decline and behavioral outcomes.

In the context of rehabilitation, emphasize the role of exercise in improving cerebrovascular health. Their findings suggest that while peripheral vascular adaptations are well-documented, the cerebral circulation's response to physical activity requires further investigation. This gap in knowledge is particularly pertinent for equine studies, where exercise is a fundamental component of health management.

[6] explore the significance of cerebral microvascular health, linking microcirculatory impairments to neurodegenerative disorders. Their review highlights the potential of MRI techniques in assessing cerebral perfusion and microvascular integrity, which could be instrumental in both human and equine medicine as a means of early detection and intervention in cerebrovascular diseases.

Lastly, [7] discuss the critical parameters of cerebral blood flow and blood volume, emphasizing the brain's energy demands and its adaptive mechanisms to maintain perfusion under varying physiological conditions. Their insights into the anatomical and functional defenses against cerebrovascular insults provide a comprehensive understanding that is applicable to both humans and horses, reinforcing the need for a comparative approach in studying cerebral circulation impairment.

This literature review aims to synthesize these findings, highlighting the shared and divergent aspects of cerebral circulation in humans and horses, and their implications for advancing both veterinary and human medical practices.

2 Methods

This research utilizes a method of comparative analysis, gathering data from veterinary and human medical studies to investigate cerebral circulation issues in both humans and horses. The analysis is centered on peer-reviewed studies and systematic reviews that address cerebrovascular health, specifically focusing on cerebral blood flow (CBF), autoregulation, neurovascular coupling, and cerebrovascular reactivity (CVR). The literature covers both experimental and clinical findings, emphasizing the similarities and differences in cerebrovascular dynamics between the two species. The reviewed studies focus on systemic hemodynamics, the anatomical structure of cerebrovascular systems, and diagnostic methods that can be applied to both humans and horses. Comparative models, particularly involving animal analogs such as pigs and horses, were examined to identify shared cerebrovascular mechanisms. The study also incorporates findings from advanced imaging techniques, like MRI, to evaluate microvascular integrity and perfusion in both pathological and physiological conditions. Diagnostic approaches were assessed for their applicability across species, with a focus on non-invasive methods for monitoring cerebrovascular health. The impact of exercise on CBF and its role in enhancing vascular health were explored, particularly in equine studies where physical activity is crucial for health management. In addition, systemic factors influencing baroreflex sensitivity and their effects on cerebral perfusion were analyzed. The review synthesizes data in order to highlight common vulnerabilities and unique adaptations in cerebrovascular systems, laying the groundwork for future interdisciplinary research. This method aims to enhance comprehension and improve strategies for managing cerebrovascular issues across different species by bringing together insights from veterinary and human medical fields.

3 Review

The article titled "Effect of propofol and remifentanil on cerebral perfusion and oxygenation in pigs: a systematic review" by [1] provides a comprehensive overview of the significance of animal models, particularly pigs, in understanding cerebral hemodynamics and the mechanisms regulating cerebral perfusion and oxygenation (CPO). The authors emphasize that the anatomical and physiological similarities between the pig brain and the human brain make pigs a valuable model for studying cerebral circulation and its implications in both veterinary and human medicine.

One of the critical insights from the article is the intricate relationship between cerebral blood flow (CBF), cerebral perfusion pressure (CPP), and cerebral vascular resistance (CVR). The authors describe how CBF is influenced by CPP, which is determined by mean arterial blood pressure (MAP) and intracranial pressure (ICP) using the formula CPP = MAP - ICP. This relationship is crucial for understanding how various physiological and pathological conditions can affect cerebral perfusion. For instance, alterations in blood pressure, cerebral metabolic rate, and other factors such as temperature and blood viscosity can significantly impact CBF, potentially leading to ischemic brain damage.

The article also highlights the importance of cerebral autoregulation, which allows the cerebral vasculature to maintain stable CBF despite fluctuations in systemic blood pressure. This intrinsic mechanism is vital for ensuring optimal cerebral metabolism and function, underscoring the need for a thorough understanding of cerebral hemodynamics in clinical settings. The authors argue that assessing cerebral perfusion and oxygenation simultaneously is essential for evaluating the physiological status of the brain and predicting clinical outcomes.

In terms of methodology, the systematic review approach taken by [1] strengthens the validity of their findings by synthesizing existing research on the effects of anesthetic agents like propofol and remifentanil on cerebral perfusion and oxygenation. This is particularly relevant for both human and veterinary medicine, as understanding the effects of these agents on cerebral circulation can inform anesthetic practices and improve patient outcomes.

The article "Human cerebrovascular function in health and disease: insights from integrative approaches" by [5] provides a comprehensive overview of the mechanisms that underpin cerebrovascular function in humans, emphasizing the evolutionary significance of these processes. The authors articulate that the brain's reliance on aerobic metabolism renders it particularly vulnerable to fluctuations in blood flow, a disadvantage that is countered by three key mechanisms: neurovascular coupling, cerebral vasoreactivity, and cerebral autoregulation.

Neurovascular coupling, as described in the article, is a critical process that ensures an adequate blood supply to active brain regions in response to metabolic demand. This coupling is essential for maintaining the brain's functionality, as it aligns regional cerebral blood flow with the specific needs of neuronal activity. The authors highlight that disruptions in neurovascular coupling can lead to neural dysfunction, indicating its pivotal role in cerebral health.

Moreover, the article titled "Impacts of systemic hemodynamic factors on cerebral and peripheral perfusion" by [3] presents a thorough investigation into the relationship between systemic hemodynamic variables and cerebral perfusion, particularly focusing on the hippocampus. The study effectively highlights the critical connection between cardiovagal baroreflex sensitivity (BRS) and regional cerebral perfusion, providing insights into how impaired BRS may correlate with cognitive dysfunction [8-9].

Laosiripisan's research underscores the significance of pulsatile blood pressure components in influencing cerebral perfusion, particularly in the hippocampus and anterior white matter. This finding is particularly relevant in the context of both human and equine medicine, as it suggests that fluctuations in blood pressure can have profound effects on brain health. The emphasis on the brain's vulnerability to variations in blood pressure is crucial, given its high-flow and low-impedance vascular characteristics. Such physiological attributes make the brain susceptible to damage from hypoperfusion, especially when steady perfusion pressure is not maintained.

Table 1. Analytical Table of Articles and Their Relevance to Veterinary and Human Medicine

Article Title	Key Insights	Relevance to Veterinary and Human Medicine
Effect of propofol and remifentanil on cerebral perfusion and oxygenation in pigs: a systematic review	- Highlights anatomical and physiological similarities between pig and human brains. - Focuses on cerebral blood flow (CBF), cerebral perfusion pressure (CPP), and cerebral vascular resistance (CVR). - Discusses the role of autoregulation in maintaining stable CBF despite systemic fluctuations.	- Provides a comparative model for understanding cerebral hemodynamics. - Informs anesthetic practices in both human and veterinary medicine.
Human cerebrovascular function in health and disease: insights from integrative approaches	- Explores cerebrovascular mechanisms such as neurovascular coupling, cerebral vasoreactivity, and autoregulation. - Emphasizes the evolutionary significance of aligning blood flow with neuronal activity.	- Explains mechanisms critical for human and equine cerebral health. - Offers insights into disease management for both species.

Impacts of systemic hemodynamic factors on cerebral and peripheral perfusion	- Investigates systemic hemodynamic factors like baroreflex sensitivity (BRS) and their impacts on regional cerebral perfusion, especially in the hippocampus. - Highlights the role of pulsatile blood pressure in influencing perfusion.	- Links cardiovascular health with cerebral perfusion. - Offers insights applicable to both human and equine physiology.
Final infarct prediction in acute ischemic stroke	- Analyzes brain energy consumption, which accounts for ~20% of energy at rest. - Discusses anatomical adaptations like the circle of Willis for maintaining perfusion. - Focuses on ischemic stroke dynamics.	- Highlights protective adaptations against cerebrovascular insults. - Applicable to both humans and animals for preventing vascular pathologies.

The article titled "Final infarct prediction in acute ischemic stroke" by [7] provides valuable insights into the dynamics of cerebral circulation, particularly in the context of ischemic stroke. The authors emphasize the critical role of the brain's energy consumption, which accounts for approximately 20% of the total energy expenditure and cardiac output at rest. This statistic underscores the importance of maintaining cerebral blood flow (CBF) and cerebral blood volume (CBV) within optimal ranges to ensure adequate delivery of oxygen and glucose to neural tissues.

A significant aspect of the article is the discussion of the brain's anatomical and functional adaptations that protect against various vascular pathologies. The authors highlight the importance of the circle of Willis, a key structure that facilitates collateral circulation, thereby playing a vital role in maintaining perfusion to the brain under compromised conditions.

4 Conclusions

The comparative analysis of cerebral circulation impairment in humans and horses reveals critical insights that have significant implications for both veterinary and human medicine. The literature reviewed establishes a foundational understanding of cerebrovascular dynamics by drawing parallels between species, particularly through anatomical and physiological similarities.

The systematic review conducted by [1] highlights the importance of maintaining stable cerebral blood flow (CBF) to prevent ischemic damage, a principle that resonates across both humans and horses. The study emphasizes the role of anesthetics on cerebral perfusion and oxygenation, providing a model for understanding CBF regulation and the intrinsic autoregulatory mechanisms that are vital for optimal cerebral metabolism.

Further, the anatomical exploration of cerebrovascular disease by [2] provides insights into the microcirculatory structures that underlie cerebrovascular health. This is complemented by the work of [3], which establishes a significant relationship between systemic hemodynamic factors and regional cerebral perfusion, particularly emphasizing the importance of baroreflex sensitivity in maintaining cerebral function.

The investigation into cerebrovascular reactivity (CVR) by [4] further underscores the shared vulnerabilities in cerebrovascular function between species. Their findings suggest that impaired CVR is linked to various neurological disorders, indicating the necessity for non-invasive assessment methods in both human and equine contexts. This is critical for advancing veterinary practices in evaluating equine neurological health.

The integrative overview provided by [5] on neurovascular coupling and its regulation in response to neuronal activity stresses the importance of understanding these mechanisms under pathological conditions. This understanding is pivotal for both species, particularly concerning cognitive decline and behavioral outcomes.

Moreover, the emphasis on exercise and its role in enhancing cerebrovascular health by points to a gap in knowledge regarding the cerebral circulation's response to physical activity, which is particularly relevant for equine studies. Lastly, the exploration of cerebral microvascular health by [6] highlights the potential for MRI techniques in assessing cerebral perfusion, which could be beneficial for early detection and intervention in cerebrovascular diseases across both fields.

In conclusion, the synthesis of findings from the reviewed articles illustrates the shared and divergent aspects of cerebral circulation impairment in humans and horses. This comparative approach not only enhances the understanding of cerebrovascular health but also underscores the necessity for integrated strategies in addressing cerebrovascular issues in both veterinary and human medical practices.

References

1. Mikkelsen, M. L. G., Ambrus, R., Miles, J. E., Poulsen, H. H., Moltke, F. B., & Eriksen, T. (2015). Effect of propofol and remifentanil on cerebral perfusion and oxygenation in pigs: a systematic review. *Acta Veterinaria Scandinavica*, *58*, 1-12.
2. Chandra, A., Li, W. A., Stone, C. R., Geng, X., & Ding, Y. (2017). The cerebral circulation and cerebrovascular disease I: Anatomy. *Brain circulation*, *3*(2), 45-56.
3. Laosiripisan, J. (2017). *Impacts of systemic hemodynamic factors on cerebral and peripheral perfusion* (Doctoral dissertation).
4. Catchlove, S. J., Pipingas, A., Hughes, M. E., & Macpherson, H. (2018). Magnetic resonance imaging for assessment of cerebrovascular reactivity and its relationship to cognition: a systematic review. *BMC neuroscience*, *19*, 1-15.
5. Ozturk, E. D., & Tan, C. O. (2018). Human cerebrovascular function in health and disease: insights from integrative approaches. *Journal of physiological anthropology*, *37*, 1-11.
6. Callewaert, B., Jones, E. A., Himmelreich, U., & Gsell, W. (2021). Non-invasive evaluation of cerebral microvasculature using pre-clinical MRI: principles, advantages and limitations. *Diagnostics*, *11*(6), 926.
7. Bertels, J., Robben, D., Vandermeulen, D., & Lemmens, R. (2022). Final infarct prediction in acute ischemic stroke. *arXiv preprint arXiv:2211.04850*.
8. Sherzod Korabayev, Jamoliddin Ergashev, Umarjon Meliboyev, Abbosbek Mukhtarov; Changes in the properties of complex pile fabrics under the influence of deformation. AIP Conf. Proc. 23 June 2023; 2789 (1): 040130. https://doi.org/10.1063/5.0145426
9. Abdurakhmon Amonov, Anvar Djuraev, Urinboy Kuryozov, Saidaposhsho Shokirova, Sherzod Korabayev; Determination of the friction force between the roller of the polymer composition coating equipment on the seams of tarpaulin materials and the surface of the tarpaulin. AIP Conf. Proc. 23 June 2023; 2789 (1): 040053. https://doi.org/10.1063/5.0145792