

# A putative mechanism of the Sodium/Iodide Symporter regulation during repetitive administration of stable Iodide described by a Systems Biology approach

David Cohen<sup>1</sup>, Dalila Lebsir<sup>1</sup>, Karine Tack<sup>1</sup>, Marc Benderitter<sup>2</sup>, and Maâmar Souidi<sup>1</sup>

<sup>1</sup> Institut de Radioprotection et de Sûreté Nucléaire (IRSN), PSE-Santé/SESANE/LRTOX, 92262 Fontenay-aux-Roses, France

<sup>2</sup> Institut de Radioprotection et de Sûreté Nucléaire (IRSN), PSE-Santé/SERAMED, 92262 Fontenay-aux-Roses, France

## Abstract

A single dose of potassium iodide (KI) against a prolonged exposure to repeated radioactivity might not be effective enough to protect the thyroid. Our group have shown that a repetitive dose of KI for eight days offers efficient protection without adverse effects in male rats [1].

However, we also have shown that the expression of the genes involved in the Wolff-Chaikoff effect changes during this period. Notably, a decrease in the sodium/iodide symporter (*nis*) gene expression has been observed [1]. This effect may result in hypothyroidism due to a decrease in thyroid hormones.

NIS is responsible for the uptake of KI and thus plays an important role in the Wolff-Chaikoff effect. The mechanism of a single dose of KI on the toxicity of the thyroid is well known [2], in contrast to repetitive administration of KI for eight days.

In the present study, we try to understand the Wolff-Chaikoff regulation and its molecular constituents during repetitive administration of KI. For this purpose, we have constructed manually a biochemical reaction network that is visualised as a “geographical” map of a single thyrocyte cell depicting the iodide and thyroid hormone metabolism. In order to investigate any regulation circuits of *nis*, Cytoscape and the plugin BiNoM [3, 4] were used to perform path analysis of the network to investigate if a path exists from the node iodide going to the node representing “*nis* transcription”. Subsequently, sequential network reduction has led to final model that might explain a putative mechanism behind *nis* regulation and repetitive iodide administration.

In addition, this map reviews the most-update information about iodide and thyroid hormone metabolism. Besides as a source of information, it can help to elucidate the mode of action of KI on gene transcription after repetitive KI administration.

## References

1. D. Lebsir, L. Manens, S. Grison, P. Lestaevel, T. Ebrahimian, D. Suhard, G. Phan, I. Dublineau, K. Tack, M. Benderitter, A. Pech, J.-R. Jourdain, M. Souidi. Mol. Cell. Endocrinol. (2018), doi: 10.1016/j.mce.2018.02.017

2. D. Carvalho, C. Dupuy. *Mol. Cell. Endocrinol.* **458**, 6 (2017)
3. P. Shannon, A. Markiel, O. Ozier, N.S. Baliga, J.T. Wang, D. Ramage, N. Amin, B. Schwikowski, T. Ideker. *Genome Res.* **13** 2498 (2003)
4. E. Bonnet, L. Calzone, D. Rovera, G. Stoll, E. Barillot, A. Zinovyev. *BMC Syst Biol.* **1** 7 (2013)