

Efficiency and genotypic quality potential of winter wheat grain in the central chernozem region

Natalia Dolgoplova^{1*}, Olga Eremenko², Gennady Feoktistov¹, and Denis Vishnevetsky¹

¹Kursk State Agrarian University named after I.I. Ivanov, 305021 Kursk, Russia

²Moscow Polytechnic University, 107023 Moscow, Russia

Abstract. Winter wheat in Russia occupies more than half of the sown areas of grain crops and more than 70% of winter crops. Advanced farms with a high culture of farming (crop rotations, fertilisers, soil treatment, etc.). The yield of this crop is more than 40 kg per ha. One of the reasons is the lack of scientifically substantiated varietal technology of cultivation of this crop in relation to soil-climatic and regional zones. As subjects of research, promising varieties, Skipetr, Moscovskaya 56, and Zorro were used. The research was conducted in 2020-2022 in the conditions of the foothill zone of the farm "Dubovitskoe" Maloarkhangel'sky district of Orel region. The paper considers the economic value of the variety, which is determined by the totality of biological and technological, food, and baking properties of high-quality varieties of agricultural crops. The choice of high-quality varieties of winter wheat and the use of the genetic resource of the variety is a decisive factor in increasing the yield of many agricultural crops, including winter wheat. When selecting the seed, it is worthwhile to assess it competently and understand all the risks of its cultivation. For many years, breeders around the world have been working on developing new productive and productive varieties. Much attention should be paid to the adaptive adaptation of varieties to external environmental conditions, which will depend on how much stable it will be possible to obtain for the programmed yield. The result of the work done shows that the new varieties under study are not inferior but superior to the control variety, which was required to be proved.

1 Introduction

The main component of increasing winter wheat yield in Russia is the selection of varieties with high adaptability to external environmental factors [1, 2]. But we should not forget that in recent years, weather conditions have been changing and many adapted varieties are not able to withstand these changes. In recent years, weather conditions have become extreme during the vegetation period of winter wheat, very often the phenomenon of drought, elevated temperatures, uncharacteristic of certain climatic zones [3, 4]. Due to this, breeders were able to identify the level of adaptability of varieties and divide them by ecological plasticity. Thus,

* Corresponding author: dunaj-natalya@yandex.ru

the selection of similar material for breeding should take into account not only the genotypic potential of grain quality, but also the reaction to changes in conditions during the period of filling and formation, the degree of its preservation in unsuitable environmental conditions [5]. Consequently, the programmes carried out by breeders should be aimed at creating adaptive forms that will be characterised by a high degree of suitability to any changes in the environment [6, 7]. It has also long been known that there is a negative correlation between quality and yield of winter wheat grain, thus, the most important task for the breeder is to develop varieties that will combine high quality and yield in new winter wheat varieties [8,9,10].

2 Research objectives

Thus, breeders are faced with the task of creating wheat varieties resistant to high plant density, with a larger reproductive part than the vegetative part, to lodging of varieties, to increase their coefficient of economic viability [11]. According to Vavilov N.I., it follows that the main way to fight against crop diseases at present is to create or select varieties that will be the most resistant to the impact of external pathogenic factors on them. In recent years, diseases caused by fungi of the genus *Fusarium* are of particular danger in winter wheat crops cultivated under intensive technology. They affect ear, grain, stem bases and root system of plants. Diseases caused by these microfungi occupy the leading place among cereal diseases by their prevalence and harmfulness [12, 13]. Significant damage to wheat grain production is caused by spikelet fusarium, which is widespread in many countries of Europe, America, Canada, and Australia.

Surveys on the study of varietal productivity of winter wheat in 2020-2022 were carried out in the circumstances of the farm "Dubovitskoe" Maloarkhangelsky district, Orel region. Full soil characteristics of the experiment location are given in Table 1.

Table 1. Soil and subsoil description of LLC Dubovitskoe, Maloarkhangelsky District, Orel region.

Characteristic features of soil	Indicators
Soil type and subtype in test plots	Black soil podzolised
Lower boundary of the rooting layer in the soil section	0.82 m
Stoniness degree	Slightly stony 2-4 m ³ /ha
Field moisture content	28.1 %
Infiltration rate	0.50-0.061 mm/min
Granulemetric composition of soils	Heavy loam
Soil volumetric weight	1.1-1.13 g/cm ³
Total soil porosity	Finely porous – 52.7 %
Humus content	5.02 %
Total nitrogen content	0.38-0.42 %
Soil acidity (pH)	Salt spray pH 5.07 Aqueous solution pH 6.18
Calcium carbonate content (CaCO ₃)	29.4 mg/eq per 100 g of soil
Base saturation	88.2-90.8 %
Mobile phosphorus content	11.02 mg/ 100 g soil
Mobile potassium content	14.83 mg / 100 g soil
Exchangeable cations composition	Ca – 29.4 % Mg – 3.3 %

In the farm winter wheat is cultivated on predecessors: black fallow, soya. The quality of seed material is shown in Table 2.

For sowing we used seeder D9-60 (12.5 cm) at the rate of 2.5 million pieces per ha.

Table 2. Seedquality

Category of seed sown	Skipetr	Zorro	Moskovskaya 56
	Original seeds	Original seeds	Original seeds
Purity %	99.80	99.57	99.00
Viability, %	98.00	96.00	99.00
Seed maturity, %	97.80	95.59	99.00

In the spring period (tillering) the following components were used:

Fungicide - Azorro 1.00 l / ha. Against brown rust, powdery mildew, leaf septoriosis, pyrenophorosis, cercosporolosis.

Herbicide - Primadonna super 0.75 litres / ha. Against annual weeds, including bush to 2.4-D and MCPA, and some perennial dicotyledons

Insecticide - Fascord 0.1-5 litres / ha. Against cereal flies, turtle bug, flea beetles, aphids, cicadas, thrips, bumblebees.

Leaf feeding - Biostim Universal 0.5 litres / ha. Ultramag combi for cereals 1.0 litres / ha.

The second treatment (2 leaves - tillering in cereal weeds)

Fungicide - Titul trio 0.6 litres / ha. Against powdery mildew, brown rust, septoriosis, pyrenophorosis, ear blight.

Herbicide - Ovsyugen express 0.5 litres / ha. Against annual cereal weeds (species of bristlewort, chicken millet, weed millet, fescue, etc.).

Insecticide - Imidor 0.1 litre / ha. Against turtle bug, intra-stem flies.

Leaf fertiliser - Ultramag combi for cereals 1.0 / ha.

The third treatment (beginning of earing)

Fungicide - Triad 0.6 litres / ha. Against Fusarium ear

Insecticide - Kinfos 0.25 l / ha. Against turtle bug, bread beetle.

Leaf fertiliser - Biostim grain 0.5 l / ha. As a system of mineral nutrition:

Main fertiliser - Diamophosca (10:26:26)

1 top dressing (beginning of tube emergence) - Ammonium nitrate (34)

2 top dressing (beginning of earing) - UAN (32).

The scheme of the experiment includes the following variants:

1 variant - variety Skipetr (test)

2 variant - variety Zorro

3 variant - variety Moskovskaya 56.

3 Methods and materials

The varieties were cultivated according to the same technology: the area of the sowing plot was 1200 m², the area of the survey plot was 100 m². Repetition in the experiment was threefold. All related analyses were carried out according to generally accepted methods and GOST. To assess the overwintering of winter wheat, the number of stems before going into winter and the number of stems after overwintering were counted. From the results obtained, the percentage of crop survival was expressed. To determine the productive bushiness, the average number of stems on one plant that yielded mature grain was calculated. The ratio of the total number of stems per 1 m² to the number of productive plants per 1 m². The length of 25 inflorescences was measured, determining the average length of the ear (from the point

of attachment of the lower ear to the top of the upper ear, not including awns) with an accuracy of 0.5 cm. The method of measuring the nature of the grain was carried out with the use of purka consists in filling with grain a measuring container with a falling weight - a measure (measuring container), having a nominal capacity of 1 dm³ and measuring the mass of this grain by weighing on scales with the subsequent exclusion from the result of weighing the mass of the empty measure with a falling weight in it. Gluten content was determined by the ratio of gluten content in the raw form to the total protein content. In our experiment we studied winter wheat varieties Skipetr and Moskovskaya 56, zoned in Region 5, as well as a promising variety of Oryol breeding Zorro.

The results obtained in the research show that technologies and effective innovative productions contribute to the solution of the main task - to obtain the highest quality products. Thus, most of the production in Dubovitsky farm is directed to seed production, which doubles the profitability of cultivated crops [14, 15]. The rest of the production is sold as marketable products - not lower than the third class. Seed products are sold in the region and abroad.

A large amount of new breeding developments of domestic and foreign science, Russian food systems and means of protection is laid down annually. That is why international conferences with the invitation of experts and specialists from the Russian Federation and friendly countries are constantly held at the Dubovitskoe base to demonstrate new technologies and agrotechnical methods of growing crops. The pilot centre has everything necessary for holding scientific seminars and conferences.

Table 3 shows the areas, production capacity and crop yields for 2020.

Table 3. Sown areas, gross production, crop yield in "Dubovitskoe" of Maloarkhangelsky district of Oryol region for 2020.

Crop	Area, ha	Yield, c/ha	Gross production, tonnes
Winter wheat	2087	65	13 566
Spring wheat	512	60	9 072
Oats	85	50	425
Peas	246	42	1 033
Buckwheat	380	22	836
Grain maize	850	85	7 225
Soybeans	1 593	25	3 983
Vica-weight	44	45	198
Millet	78	30	234
Sugar beet	30	600	1 800
Forage crops and other	425	-	-
Total	7 330		

4 Results

Data on overwintering of winter wheat varieties are presented in Table 5.

According to the data of Table 4, it follows that variety Zorro best passed the overwintering stage (95%), which is more than the control (Skipetr) by 17%. The Moskovskaya 56 variety had 2 % lower overwintering readings than the control.

Table 4. Effect of variety peculiarities on overwintering of cereals

Experimental sample	Number of stems in autumn, pcs./1 plant	Number of stems in spring, pcs./1 plant	Pre-winterisation process, %
Skipetr (test)	3.6	2.8	78
Zorro	3.8	3.6	95
Moskovskaya 56	3.7	2.8	76

The results of the conducted research show that the studied varieties of cereals have a tangible effect on the indicators of elements of yield structure (Table 5).

The highest 1000 grains of winter wheat is achieved by variety Zorro - 41.1 g, which is 4.2 g higher than the standard Skipetr.

Table 5. Variety influence on elements of winter wheat yield structure

Wheat variety	Productivity in plant bushiness	Spike length, cm	Number of grains per ear, pcs.	Grain weight in ear, g	Weight of 1000 grains, g
Skipetr (test)	2.1	8.6	30	1.10	36.9
Zorro	2.3	9.5	38	1.12	41.1
Moskovskaya 56	2.2	7.4	27	1.06	34.2

The highest coefficient of productive bushiness of winter wheat was provided by variety Moskovskaya 56 - 2.2, which is higher than the standard Skipetr by 0.1 and variety Zorro (+0.2 to control). Variety Zorro showed better indicators for ear length, grain weight in the ear and grain in it (9.5 cm, 1.12 g, 38 pcs). That is higher than the control by 0.9 cm, 0.02 g and 8 pcs of grains, respectively. The yield of winter wheat was influenced by weather conditions and genetic features of cultivated varieties. Thus, under the conditions of 2020 this indicator by varieties ranged from 51.0 kg / ha for variety Moskovskaya 56 to 78.6 kg / ha for variety Zorro.

Table 6. Yield of winter wheat varieties in the farm "Dubovitskoe" of Maloarkhangelsky district of Orel region

Winter wheat variety	Yield, c/ha	Deviation from control, +/-	
		c/ha	%
Skipetr (test)	71.5	-	-
Zorro	78.6	+7.1	+ 9.9 %
Moskovskaya 56	51.0	-20.4	- 28.5 %
HCP ₀₅		4.8	

In the Zorro variety, yields increased by 7.1 kg/ha, or 9.9 %. At the same time, the yield of Moskovskaya 56 decreased by 20.4 kg/ha (28.5 %)

Researches have established that winter wheat varieties provide gluten content in grain at the level of 23.9 - 26.6 % (Table 7).

Table 7. Technological qualities of winter wheat grain depending on variety

Winter wheat variety	Natura, g/l	Vitreousness, %	Gluten, %	Gluten quality, units. FDM	Quality Group
Skipetr (test)	775	45	23.9	77	II
Zorro	765	47	26.6	85	II
Moskovskaya 56	725	44	25.9	84	II

The quality and quantity of gluten in grain is a genetically inherited trait. As a result, the highest quality of gluten is achieved in the grain of winter wheat variety Zorro is 26.6 %, which is higher than Skipetr variety by 2.7%. According to the data of cultivation of the studied varieties observed that the highest yield was Zorro variety 73.8 kg / ha, which is higher than the control (Skipetr variety) by 13 kg / ha.

5 Conclusions

Based on the research conducted in 2020-2022 in the conditions of the farm "Dubovitskoe" of Maloarkhangelsky district of Orel region to study the varietal productivity of winter wheat, the following conclusions can be drawn:

- The variety has a significant effect on the level of winter wheat overwintering: in the control variety, Skipetr overwintering was 78 %, and in the Zorro variety increased by 17 %, reaching the level of 95 %.
- Compared to the control, the yield of Zorro variety increased by 7.1 kg/ha, or 9.9 %.
- The best indicators of the technological qualities of winter wheat grain were observed in the Zorro variety: vitreousness increased by 2 %, crude gluten content - by 2.7 %.
- The maximum grain texture was formed on the control variety Skipetr and was 775 g/l, which is higher than that of Zorro variety by 10 g/l and Moskovskaya 56 variety - by 50 g/l.

References

1. A.V. Gostev, *Zemledeliye* **6**, 16-20 (2019) <https://doi.org/10.24411/0044-3913-2019-10604>
2. I.A. Goneev, O.P. Lukashova, E.A. Batrachenko, V.N. Lunin, *Problems of regional ecology* **2**, 56-63 (2020) <https://doi.org/10.24411/1728-323X-2020-12056>
3. E.N. Zubkova, T.A. Belova, *Bulletin of Peoples' Friendship University of Russia. Series: Ecology and life safety* **25(1)**, 50-57 (2017) <https://doi.org/10.22363/2313-2310-2017-25-1-50-57>
4. S.D. Karakotov, A.I. Pryanishnikov, et al. *Bulletin of Agrarian Science* **3(102)**, 7-17 (2023) <https://doi.org/10.17238/issn2587-666X.2023.3.7>
5. E.A. Batrachenko, N.V. Dolgopolova, T.A. Dudkina, *Investigation of the soil cover ecological state under the different geomorphological elements' conditions*. In Proceedings: IOP Conference Series: Earth and Environmental Science. Krasnoyarsk Science and Technology City Hall. Krasnoyarsk, Russian Federation, 2021. P. 42081.
6. E.A. Batrachenko, I.A. Goneev, O.P. Lukashova, V.V. Rudnev, *Diagnostics of the ecological state of aquatic and urban ecosystems using radiometric and bioindication analysis methods*. In Proceedings: JOP Conference Series: Metrological Support of Innovative Technologies. Krasnoyarsk Science and Technology City Hall of the

- Russian Union of Scientific and Engineering Associations. Krasnoyarsk, Russia, 2020. P. 32066. <https://doi.org/10.1088/1742-6596/1515/3/032066>
7. T.A. Dudkina, *Tavricheskiy vestnik agrarnoy nauki* **1**, 30-40 (2022) <https://doi.org/10.33952/2542-0720-2022-1-29-30-40>.
 8. N.V. Dolgoplova, I.Ya. Pigorev, *Bulletin of Kursk State Agricultural Academy* **6**, 7-11 (2017)
 9. T.A. Dudkina, *Bulletin of the Ulyanovsk State Agricultural Academy* **4**, 14-20 (2022) <https://doi.org/10.18286/1816-4501-2022-4-14-20>.
 10. I.G. Pykhtin, A.V. Gostev, A.I. Pykhtin, *Journal of Engineering and Applied Sciences* **12(20)**, 5338-5342 (2017) <https://doi.org/10.3923/jeasci.2017.5338.5342>
 11. I.Y. Pigorev, O.V. Nikitina, *Bulletin of the Ryazan State Agrotechnological University named after P.A. Kostychev* **15(2)**, 45-51 (2023) <https://doi.org/10.36508/RSATU.2023.84.12.007>
 12. S.V. Rezvyakova, V.N. Titov, et al. *Agrarnayanauka* **2**, 76-81 (2023) <https://doi.org/10.32634/0869-8155-2023-367-2-76-81>
 13. A.V. Gostev, *Feeding of agricultural animals and fodder production* **2(211)**, 47-71 (2023) <https://doi.org/10.53859/02352451-2022-26-8-5>
 14. N.V. Dolgoplova, A.S. Akimenko, *Bulletin of the Kursk State Agricultural Academy* **1**, 11-15 (2017)
 15. A.I. Stifeev, O.V. Nikitina, *Bulletin of the Kursk State Agricultural Academy* **2**, 30-35 (2022)