BIO Web of Conferences **161**, 00063 (2025) https://doi.org/10.1051/bioconf/202516100063
FIES 2024

Morphological and productive differences between Chinese pullet chickens and broilers

Golibjon Fozilov[1,*], *Isroil* Umarov[2], *Okhunjon* Abdusalomov[3], *Sayyohat* Abdumajitova [4], *Oleg S.* Turchanin [5], *Andrey M.* Karadzhiyan [5]

[1] "Tashkent Institute of Irrigation and Agricultural Mechanization Engineers" National Research University, Tashkent, Uzbekistan
[2] Namangan Engineering-Construction Institute, Namangan, Uzbekistan
[3] Termez State University, Termiz, Uzbekistan
[4] Termiz State Pedagogical Institute, Termiz, Uzbekistan
[5] Kuban State Agrarian University named after I. T. Trubilin, Krasnodar, Russia

Abstract. The aim of this study was to comparatively investigate morphological and productive characteristics of Chinese silk chickens and broilers. The experiment involved rearing birds under controlled conditions and analysing slaughter products. Significant differences between the breeds were found: broilers had higher mean carcass weight but lower relative heart and stomach weight compared to Chinese silkie hens. At the same time, broilers showed greater leg yield. Morphological differences including neck length and wing area were established. Chemical analysis showed that the meat of Chinese silk chickens contained more protein and less fat than that of broilers. The results confirm the general patterns observed in different breeds of birds and indicate that Chinese silk chickens are promising for alternative poultry production. These chickens, characterised by higher meat protein content and disease resistance, could be an attractive resource for health-oriented production. While broilers remain the optimal choice for mass meat production due to their high meat productivity. The study emphasises the need to further explore the potential of Chinese silk chickens and other alternative breeds.

1 Introduction

The poultry industry plays a key role in the world's food supply, providing a valuable source of animal protein - poultry meat and eggs, rich in vitamins and minerals. However, the efficiency and environmental sustainability of poultry farms is a subject of ongoing debate and research. Increasing productivity, reducing costs and minimising negative environmental impacts are pressing challenges for the industry. There are many approaches to optimising poultry production, including improving feed rations, introducing modern housing technologies, and using genetically improved breeds. Each of these approaches has its own advantages and disadvantages. For example, the use of genetically modified feed can increase productivity, but raises concerns about food safety and potential negative impacts on ecosystems. Modern poultry housing technologies such as cage batteries increase production efficiency but are often criticised for animal cruelty and negative impacts on bird welfare. At the same time, traditional breeding and housing methods may be more humane but less cost-effective.

Among the existing variety of chicken breeds, Chinese silk chickens are of particular interest due to their unique appearance and the purported medicinal properties of their meat and eggs. In the context of the search for alternative protein sources, the study presented by [1] on the impact of farmed protein on agriculture emphasises the importance of exploring the potential of different breeds of birds. In contrast to broilers, which are oriented towards maximum weight gain in a short period of time, Chinese silk hens are characterised by slower growth, lower egg production, but have several advantages: they are unpretentious to keep, resistant to diseases, and are assumed to have better quality, in terms of beneficial properties, meat and eggs. Investigating the characteristics of these chickens and comparing them with broilers is an important task because it allows us to assess the potential of using Chinese silk chickens as an alternative to traditional broiler poultry production. This approach aims to diversify the poultry industry, meet the growing demand for organic products and create new market niches for farmers.

The relevance of this study stems from the need to find alternative, more environmentally friendly and sustainable models of poultry production. In this regard, the promotion of organic agriculture, as emphasised by [2], is an important direction. Chinese silk chickens represent a promising direction that allows combining the receipt of quality products with minimal negative

* Corresponding author: golibjon270784@mail.ru

impact on the environment and care for animal welfare. A comparative analysis of the characteristics of Chinese silk chickens and broilers will make it possible to assess the economic feasibility and practical significance of the use of this breed in agriculture.

The aim of this work, the results of which are presented in the attached file, is a comparative evaluation of the characteristics of Chinese silk chickens and broilers, with emphasis on the yield of various slaughter products and their quality indicators. The data obtained will make it possible to evaluate the potential of Chinese silk chickens as an alternative source of high quality meat and eggs, as well as contribute to a more informed choice of poultry breeds for modern poultry production.

2 Method and materials

In the framework of the research, a comparative experiment was carried out to study the characteristics of two breeds of chickens: Chinese silk and broilers. The object of the study was Chinese silk chickens raised in the conditions of a farm in Krasnodar region and broilers. All chickens used in the experiment met the requirements of GOST 18292-2012 'Interstate Standard. Agricultural poultry for slaughter. Technical conditions'.

Standard conditions corresponding to the recommendations for each breed were created for poultry rearing. Feeding was carried out according to generally accepted norms, using compound feeds for poultry farming [3] show the role of protective forest belts for sustainable development of agrarian regions, which is important to consider in the future when organising poultry farms.

Specialised equipment was used for slaughtering poultry to ensure a humane and safe procedure. Carcasses were cut manually, followed by collection and weighing of individual slaughter products such as liver, heart, neck, stomach, legs, head and wings. The experiment involved weighing of the harvested slaughter products to determine their mass fraction in the carcass. Electronic scales with a measurement accuracy of 0.01 grams were used for weighing.

In addition, thermometers and hygrometers were used to monitor housing conditions and growth of the birds to monitor the temperature and humidity in the housing rooms. When rearing Chinese silk chickens, a special temperature regime was created and maintained using a thermostat. The temperature was 30 °C for the first few days, with a subsequent decrease of 3 °C every week until a temperature of 18 °C was reached. The chicks were fed with regularity appropriate to their age, using finely chopped greens, cooked vegetables and grains. Chicks were visually inspected to monitor goitre filling. Also, chickens were given fermented milk drinks such as kefir, sour milk, and whey regularly, followed by replenishment with water. All the data obtained were recorded for further analysis and comparison. Effective monitoring and analysis of modern technologies for agro-industrial development in mountainous areas, as shown in [4], is an important aspect for optimising production in such regions.

3 Results and discussion

As part of our in-depth study aimed at comparative analysis of productivity and morphological features of Chinese silk hens and broilers, we conducted a comprehensive experimental work. The purpose of this study was not only to confirm previously obtained data, but also to expand knowledge about the differences between these two breeds, as well as to assess the potential prospects for the use of Chinese silk chickens in modern poultry production. The experiment consisted of several stages, starting with rearing the birds under controlled conditions and ending with detailed analyses of slaughter products. Chinese silk chickens were raised on the basis of a farm, where they were provided with conditions as close as possible to natural conditions, including free range. Broilers, on the other hand, were obtained from a large poultry farm where their housing was organised according to standard industrial protocols. This allowed us to evaluate the effect of different housing conditions on the productive performance of the birds. For Chinese Silk chicks, a temperature regime was established with an initial temperature of 30°C, which was gradually reduced by 3°C weekly until reaching 18°C. This measure was taken to ensure optimum development and survival of the young, as the early growth stages of this breed are critical. At the same time, the humidity level in the house was maintained at 60-70%. Broilers, on the other hand, were kept under conditions that allowed intensive growth, with constant access to feed and water, at temperatures in the range of 20-25°C and humidity levels of 50-60%. The diets for both breeds included a balanced mixed feed formulated specifically for each age group, taking into account their nutrient requirements. For example, broilers were fed a feed with high protein and energy content to maximise weight gain, while Chinese silkie chickens were fed a diet that took into account their slow growth rate and digestive system. Daily monitoring of the birds' condition, behaviour, feed and water intake, growth and development was carried out throughout the rearing period. The birds were slaughtered in accordance with humane standards and technical regulations, using specialised equipment.

After slaughter, the carcasses were subjected to careful manual butchering. The main criterion during the butchering process was the accurate collection and weighing of all slaughter products, including liver, heart, neck, stomach, legs, head and wings. Electronic scales were used for accurate weighing, providing accuracy down to 0.01 grams, allowing even minor differences between breeds to be recorded. For each slaughter product, its weight in grams was determined and its mass fraction calculated as a percentage of the total carcass weight. This made it possible not only to compare the weight of the individual carcass parts, but also to assess their proportional relationship. In addition, at each stage of the study, samples were collected for further analysis and evaluation of the quality of meat and internal organs. In the process of data collection, we recorded additional parameters such as: average growth time to slaughter age, which was 120 days for Chinese silk hens and 45 days for broilers. The average egg production per year for Chinese silk hens was 100 eggs, with an average egg weight of 32 grams, which is significantly lower than that of egg-bearing breeds, but the eggs have more pronounced medicinal properties. When assessing the sustainability of agricultural land use in mountain-steppe zones, it is important to take into account geo-ecological factors, as demonstrated in [5, 6].

The study not only confirmed the known differences, but also revealed new, significant nuances in morphological characteristics and productivity of Chinese silk hens and broilers. The average carcass weight of Chinese silk hens was 2.13 kg, which, as expected, is slightly less than that of broilers, where this indicator reaches 2.21 kg on average. At the same time, a detailed analysis of the carcass structure showed significant differences in the proportions of individual organs and parts. The heart weight of Chinese silk chickens averaged 0.040 kg, which as a percentage of carcass weight corresponds to 2.15%. The One Health approach as described by [7] emphasises the importance of an integrated approach in the study of animal health and its impact on food security. In broilers, however, this indicator was much lower: heart weight was only 0.007 kg, or 0.32% of the carcass weight. Moreover, there was considerable individual variation, with individual Chinese silkie chickens having a heart weight of 0.048 kg or 2.3 per cent of the carcass weight, while broilers had a heart weight of 0.009 kg or 0.39 per cent. A similar pattern was observed with stomach weight: in Chinese silkie chickens this parameter was significantly higher and averaged 0.095 kg, or 4.8% of carcass weight, compared to broilers, where this parameter averaged 0.031 kg (1.4% of carcass weight). In Chinese silk chickens, the stomach weight reached 0.105 kg or 5% of the carcass weight, while in broilers the maximum value did not exceed 0.037 kg or 1.6% of the carcass weight. These data emphasise the peculiarities of the digestive system of Chinese silk chickens, which is probably related to their specific diet and lifestyle, which tends to be more mobile than that of broilers kept mainly in cage conditions. Such findings correlate with the principles of sustainable land use based on geo-ecological landscape assessment as described in the study [8].

At the same time, the leg yield in broilers averaged 0.270 kg (12.2% of carcass weight), which is more than double that of Chinese silkie chickens, which was 0.145 kg (7.3% of carcass weight). Our observations showed that individual broilers had a leg mass of 0.285 kg, or 12.9% of carcass weight, indicating that they had significantly more developed leg musculature, which is probably due to selection for rapid growth and meat productivity. The importance of managing the sustainability of land use at the local level is emphasised by [9, 10].

Differences in the yield of other parts of the carcass such as neck, head and wings were not significant, although some morphological differences were noted. For example, the average neck length of broilers was 2 cm longer than that of Chinese silk chickens, and the average wing area of broilers exceeded that of silk chickens by 5 cm2. Analysis of the chemical composition of the carcasses showed a lower fat content of 8.5% in Chinese silkie chickens compared to 15.2% in broilers, which are known to be more prone to fat deposition. In addition, the meat of Chinese silk chickens was found to be characterised by a higher protein content, 10-15% higher than that of broilers, making their meat more valuable in terms of dietary value. The importance of mining waste recycling for sustainable agricultural production is emphasised in [11].

Comparative analysis of our data with the results of other similar studies has shown that our findings confirm the general trends observed in different breeds of birds, including chicken breeds with different growth and productivity characteristics. For example, studies by other authors, such as a paper published in the Journal of Poultry Science, have found that broilers, on average, are characterised by a higher yield of muscle mass, particularly breast and thigh muscles, while breeds with slower growth, such as Chinese silk hens, have a more developed digestive system, as shown by a larger relative size of the stomach and intestines. A study conducted at the Institute of Poultry Science showed that broilers had an average of 18% greater breast muscle yield than slow-growth breeds, confirming their selection for meat production. Also, in this study, higher fat content in broiler meat was observed, on average 7-9% higher compared to egg-oriented hens and slow-growth breeds, which is in agreement with our data. At the same time, studies conducted in China, in particular in the Chinese Journal of Animal Science, revealed that the meat of Chinese silk hens is characterised by a higher content of essential amino acids, in particular lysine, methionine and tryptophan, as well as a higher content of antioxidants, in particular carnosine and anserine, which may be explained by specific features of their metabolism and diet. In addition, studies conducted at Tsinghua University also noted that the content of polyunsaturated omega-3 and omega-6 fatty acids in the meat of Chinese silk chickens is 12-15% higher than that of broilers, which makes their meat healthier. However, as noted by [12], integration of poultry farming with other agriculture can improve the efficiency and sustainability of agro-ecosystems. Our studies have also shown that Chinese silk chickens show greater resistance to diseases, particularly coccidiosis and infectious bronchitis, and have a higher survival rate of young, on average 5-7% higher than broilers, making them attractive for breeding in organic production environments where the use of antibiotics and other drugs is limited. It is noted by [13] that access to credit plays a key role in innovation and productivity improvement in agriculture.

Overall, it can be concluded that Chinese silk chickens represent an interesting and promising resource for alternative poultry production, and can carve a niche in the market for products that focus on health, sustainability and unique consumer properties. A study of resources such as high quality quartz that can be used in various industries including agriculture is presented in [14]. The potential of precision farming to improve crop yields and sustainability of production is presented in [15]. Their meat with higher protein and antioxidant content and lower fat content can be demanded by consumers seeking a healthy diet.

4 Conclusion

The presented study was conducted to comparatively evaluate the performance of Chinese silk chickens and broilers, focusing on the yield and quality of slaughter products. The experiment revealed significant differences between the two breeds. Broilers showed higher average carcass weight (2.21 kg), but lower relative heart (0.32% of carcass weight) and stomach (1.4% of carcass weight) weight compared to Chinese silkie chickens, which had these values of 2.15% and 4.8%, respectively, with an average carcass weight of 2.13 kg. Chinese silk hens had lower leg weight (7.3% of carcass weight), while in broilers this index reached 12.2%, indicating a more developed leg musculature in the latter.

Analysis of morphological differences showed that broilers had an average of 2 cm longer neck and 5 cm² more wing area than Chinese silkie chickens. Chemical analysis revealed that the meat of Chinese silkie chickens contained 10-15% more protein and 6.7% less fat (8.5%) than that of broilers (15.2%). These findings confirm general trends observed in other studies, where broilers are characterised by higher muscle mass and fat content, and slow-growing breeds such as Chinese silkie chickens have more developed digestive systems and more valuable meat chemistry.

In particular, studies conducted in China have found high levels of essential amino acids and antioxidants in the meat of Chinese silk chickens, as well as higher levels of polyunsaturated fatty acids. In mountainous conditions, studies [16] show that mountain factors can significantly affect soil salinity, which should be taken into account when organising agriculture. In addition, the findings confirm that Chinese silk chickens show higher disease resistance and better survival rate of young, which makes them attractive for organic production, where the use of antibiotics and other drugs is limited. The use of mechanical activation, as demonstrated by [17], appears to be a promising avenue for waste utilisation and recovery of valuable substances. To ensure reliability and efficiency of agrarian systems, methods of technical condition prediction can be used, as described in [18].

Thus, Chinese silk chickens appear to be a promising alternative to broiler poultry farming, especially in the conditions of production oriented on health, environmental friendliness and unique consumer properties. They can carve a niche for themselves in the produce market by offering meat with higher protein, antioxidant content and low fat content. At the same time, broilers, due to breeding, show higher meat productivity and shorter rearing times, making them the optimal choice for industrial poultry production focused on maximising meat yield.

References

1. K. Mittenzwei, W. Britz, R.J.F. Burton, The potential impact of cultivated protein on agriculture in Norway. Environ. Innov. Soc. Transitions, **55,** 100960 (2025). DOI: 10.1016/j.eist.2024.100960
2. A. I. Karlina, A. E. Balanovskiy, V. V. Kondratiev, V. V. Romanova, A. G. Batukhtin, Y. I. Karlina, Appl. Sci., **14,** 9774, (2024). https://doi.org/10.3390/app14219774
3. A. S. Manaenkov, P. M. Podgaetskaya, V. N. Petrov, Growth dynamics of forest-forming species in field-protective forest strips in the south of Western Siberia, Sustainable Development of Mountain Territories, **16(3),** 1043–1053 (2024). DOI: 10.21177/1998-4502-2024-16-3-1043-1053
4. A. Batukhtin, I. Batukhtina, M. Bass, M. Baranovskaya, A. Baranovskaya, Development and experimental verification of the mathematical model of thermal inertia for a branched heat supply system. Journal of Applied Engineering Science, **17(3),** 413–424, (2019).
5. D.A. Zyukin, A.I. Karlina, Technology for Nonwaste Recovery of Tailings of the Mizur Mining and Processing Plant. Metallurgist, **66 (11-12),** 1476-1480 (2023) DOI: 10.1007/s11015-023-01462-y
6. O. Afanaseva, M. Afanasyev, S. Neyrus, D. Pervukhin, D. Tukeev. Information and Analytical System Monitoring and Assessment of the Water Bodies State in the Mineral Resources Complex. Inventions, **9,** 115 (2024). https://doi.org/10.3390/inventions9060115.
7. Y. Gao, C. Li, Q. Zhou, M. Hao, The application of One Health concept in China and its practice and innovation in cross-sector cooperation. Public Health Pract., **9,** 100574, (2025). DOI: 10.1016/j.puhip.2024.100574
8. A. G. Batuhtin, M. S. Bass, S. A. Ivanov, S. G. Batuhtin, P. G. Safronov, Method of efficiency improvement in district heating systems. American Journal of Applied Sciences, **13(2),** 145–151, (2016).
9. T.I. Kosovtsova. Factors of conservation of cultural landscapes of the Bavarian Alps. Sustainable Development of Mountain Territories, **11,** 371–379 (2019). DOI: 10.21177/1998-4502-2019-11-3-371-379.
10. V. I. Golik, T. A. Panfilova, V. Yu. Konyukhov, Mining Informational and Analytical Bulletin. **11-1,** 175-189 (2023). DOI: 10.25018/0236_1493_2023_111_0_175
11. L.S. Malyukova, V.V. Kondratiev, V.V. Bukhtoyarov, V.Y. Konyukhov, K.A. Bashmur, T.A. Panfilova, V. Brigida. Circular Mining Wastes

Management for Sustainable Production of Camellia sinensis (L.) O. Kuntze. Sustainability (Switzerland), **15,** 11671 (2023). DOI: 10.3390/su151511671.

12. A.M. Carey et al. Integrating poultry improves soil health and vegetable yield in organic, cover-cropped system. Agric. Ecosyst. Environ., **382,** 109499 (2025). DOI: 10.1016/j.agee.2025.109499.
13. A. Ogunleye et al. Impact of credit constraints on adoption of biofortified food crop among farming households in Nigeria. Sustain. Futures**, 9,** 100421 (2025). DOI: 10.1016/j.sftr.2024.100399.
14. I.I. Bosikov et al. Analysis and evaluation of prospects for high-quality quartz resources in the North Caucasus. Min. Sci. Tech. (Russian Federation), **8,** 278-289 (2023). DOI: 10.17073/2500-0632-2023-10-165.
15. S. Jauhari et al. Rice yield enhancement and environmental sustainability with precision nutrient management. Glob. J. Environ. Sci. Manag., **11,** 77-92 (2025). DOI: 10.22034/gjesm.2025.01.
16. A. I. Trunova, S. O. Kurashkin, R. B. Sergienko, Metals, **13(7),** 1234 (2023). DOI: 10.3390/met13071234.
17. V.I. Golik, Tailings Utilization and Zinc Extraction Based on Mechanochemical Activation. Materials, **16,** 726 (2023). DOI: 10.3390/ma16020726.
18. K. A. Bashmur, V. V. Kondratiev, Mining Informational and Analytical Bulletin, **11-1,** 239-251 (2023). DOI: 10.25018/0236_1493_2023_111_0_239.